

Debye Potentials for Maxwell and Dirac Fields
from a
Generalisation of the Killing-Yano Equation.

I. M. Benn¹
Philip Charlton²
Jonathan Kress³

Mathematics Department,
Newcastle University,
NSW 2308,
Australia.

Abstract. By using conformal Killing-Yano tensors, and their generalisations, we obtain scalar potentials for both the source-free Maxwell and massless Dirac equations. For each of these equations we construct, from conformal Killing-Yano tensors, symmetry operators that map any solution to another.

¹ email:mmimb@cc.newcastle.edu.au
² email:jkress@maths.newcastle.edu.au
³ email:philipc@maths.newcastle.edu.au

I. Introduction.

Maxwell's equations require that the Maxwell 2-forms be closed, and hence locally exact. They are globally exact if we discount the existence of magnetic monopoles, and Maxwell's equations can then be written as second-order equations for a potential 1-form. In the absence of sources the Maxwell 2-forms are also co-closed and hence locally co-exact. For charge-free solutions they are globally co-exact and a potential 3-form can be introduced. A Hertz potential is a 2-form in terms of which the Maxwell 2-form can be expressed so as to be simultaneously exact and co-exact, and hence satisfy the source-free Maxwell equations. In certain cases one can parameterise the Hertz potential in terms of a function satisfying a second-order differential equation. Thus Maxwell solutions can be expressed in terms of a scalar potential: the Debye potential. (Clearly we have paraphrased what Hertz and Debye actually did. Although we shall not attempt to give an historical account it might be noted that Bromwich and Whittaker also contributed to these ideas. We have given references later.)

Some of the important solutions to Maxwell's equations in flat space are expressed in terms of solutions to scalar equations. If the Lorenz gauge condition is imposed on the potential 1-form then Maxwell's equations require that it be harmonic. In a parallel basis this requires that its components be harmonic functions. The plane wave solutions may be obtained in this way. The radiating multipole solutions are adapted to spherical symmetry. Here the electric and magnetic fields are required to satisfy a vector Helmholtz equation. Solutions to this vector equation can be expressed in terms of solutions to a scalar Helmholtz equation. Thus the radiating multipole family can be expressed in terms of scalar potentials. In a curved space-time one cannot immediately generalise these solutions, as the derivatives introduce extra connection terms. Cohen and Kegeles [1] seem to have been the first to apply the Hertz potential formalism to Maxwell's vacuum equations in a curved background. They point out that the Hertz potential must be some geometrically privileged 2-form, and that algebraically-special space-times have such 2-forms, corresponding to the repeated principal null directions.

Sections II to V are all preliminaries to the Debye potential formalism. A preliminary reading may begin at section VI, referring to the earlier sections as necessary. Section II introduces some notation, whilst Section III expresses some results on the Petrov classification scheme in a form that will be convenient later. In Section IV we introduce the conformal Killing-Yano equation. This conformal generalisation of Yano's Killing-tensor equation was introduced in [2]. We show how the equation can be written equivalently in terms of exterior operations. This provides an elegant statement of the equation. Moreover, it is natural to have the equation expressed in this way when considering Maxwell's equations which are themselves most naturally expressed in exterior form. Any differential form may be regarded as a tensor on the spinor space. We show how the conformal Killing-Yano equation for a 2-form is equivalent to an equation for a 2-index Killing spinor.

This relationship has not always been made clear: indeed, tensors equivalent to Killing spinors have been called Penrose-Floyd tensors. In Section V we consider equations for self-dual 2-forms whose eigenvectors are shear-free. When the 2-forms have only one real eigenvector then Robinson's theorem [3] says that they are proportional to a closed form. We show that equivalently the 2-forms satisfy a 'gauged' conformal Killing-Yano equation. This restatement of Robinson's theorem proves convenient for obtaining Debye potentials. When the 2-form has two real shear-free eigenvectors then we recover the equations obtained by Dietz and Rüdiger [4]. We obtain the integrability conditions for these shear-free equations that we will need later. We do this by relating the self-dual 2-forms to spinor fields. This is not only a convenient way of obtaining these integrability conditions, but also enables us to consider the Debye scheme for the Dirac equation in section VII.

In section VI we show how Debye potentials for Maxwell fields are related to repeated principal null directions in algebraically-special space-times. This had previously been done in Newman-Penrose formalism by Cohen and Kegeles [1, 5]. Thus their Debye potential equations were written out explicitly in an adapted basis. The advantage of our approach is that we obtain the equations in a basis-independent way, which we feel makes it easier to see how the various ingredients to the scheme enter. (Of course, to solve the equations in any given space-time necessitates choosing some basis adapted to the geometry.) Other workers have utilised different aspects of special properties of space-times admitting Debye potentials. Stewart [6] considered Petrov type D vacuum space-times and used the more specialised Geroch-Held-Penrose formalism to obtain simplified equations. Torres del Castillo [7] emphasises the special properties of totally null foliations or 'null strings' in his treatment of null Hertz potentials. Wald [8] has pointed out the nature of the relationship between Cohen and Kegeles' Debye potential equations the decoupled massless field components of Teukolsky [9, 10] from which he derives the Debye potential equations.

In the special case in which there exists a conformal Killing-Yano tensor the Debye potential scheme becomes both simpler and more powerful. (Here our way of writing the conformal Killing-Yano equation is particularly well adapted to the scheme.) It is possible to use Debye potentials to obtain a symmetry operator for Maxwell's equations, mapping any source-free Maxwell solution to another. This has been shown by Torres del Castillo [11] using 2-index Killing spinors. Now Kalnins, McLenaghan and Williams [12] have obtained the most general second order symmetry operator for the source-free Maxwell system. Their operator contains a term constructed from a 4-index Killing spinor. We show how such an operator is obtained from the Debye potential scheme. We show how it is expressed in terms of a 'generalised conformal Killing-Yano tensor'.

In section VII we treat the massless Dirac system analogously to the Maxwell system. We show the relation to previous work, and in particular show how the Debye potential method generalises Penrose's 'spin raising' and 'spin lowering' operators, constructed from twistors in conformally-flat space [13], to an algebraically-special space-time.

In the final section we summarise our results and discuss possible generalisations.

II. Notation and Conventions.

The exterior calculus of differential forms will be used extensively. As usual $\wedge$ denotes the exterior product and d the exterior derivative. If X is a vector field then $X\lrcorner$ denotes the interior derivative that contracts a p -form with X to produce a $(p-1)$ -form. The interior derivative is an anti-derivation on differential forms (as is d) with $X\lrcorner A = A(X)$ for A a 1-form. It follows that if ω is a p -form then

$$e^a \wedge X_a \lrcorner \omega = p\omega$$

where the coframe $\{e^a\}$ is dual to the frame $\{X_a\}$.

On a pseudo-Riemannian manifold the metric tensor g establishes a natural isomorphism between vector fields and differential 1-forms. The 1-form $X^\flat$ is defined such that $X^\flat(Y) = g(X, Y)$ for all vector fields Y . Thus $X^\flat$ has components obtained by lowering those of X with the metric tensor. The inverse of $\flat$ is $\sharp$, resulting in the vector $\omega^\sharp$ having components obtained by raising those of the 1-form ω .

The metric tensor also gives a natural isomorphism between the space of p -forms and the $(n-p)$ -forms, where n is the dimension of the manifold. This isomorphism is the Hodge duality map $*$. On a decomposable p -form we have

$$*(X_1^\flat \wedge X_2^\flat \wedge \dots \wedge X_p^\flat) = X_p \lrcorner X_{p-1} \lrcorner \dots X_1 \lrcorner *1 \quad (1)$$

where $*1$ is the orienting volume n -form. *We shall only consider four-dimensional Lorentzian manifolds.* The metric tensor g will be taken to be positive-definite on space-like vectors. With these conventions we have

$$**\omega = \begin{cases} \omega & \text{if } \omega \text{ is of odd degree;} \\ -\omega & \text{if } \omega \text{ is even.} \end{cases} \quad (2)$$

The Hodge dual may be combined with the exterior derivative to form the co-derivative d^* that acts on a form to lower the degree by one. For the case of four dimensions and Lorentzian signature

$$d^* = *d* . \quad (3)$$

The exterior derivative and the co-derivative can be expressed in terms of the Lorentzian connection ∇ by

$$d = e^a \wedge \nabla_{X_a}$$

and

$$d^* = -X^a \lrcorner \nabla_{X_a} .$$

The Clifford algebra of each cotangent space is generated by the basis 1-forms. It may be identified with the vector space of exterior forms with Clifford multiplication related to the exterior and interior derivatives by

$$A\phi = A \wedge \phi + A^\sharp \lrcorner \phi$$

for A a 1-form and ϕ an arbitrary form. We will juxtapose symbols to denote their Clifford product. Whereas the Clifford product of two homogeneous forms will be inhomogeneous, the Clifford commutator of a 2-form with another form will preserve the degree of that form. The Clifford commutator, denoted $[\ , \]$, is related to the exterior product by

$$[F, \phi] = -2X_a \lrcorner F \wedge X^a \lrcorner \phi$$

where F is a 2-form and ϕ is an arbitrary form. The Lie algebra formed by the 2-forms under Clifford commutation is the Lie algebra of the Lorentz group. (In general they form the Lie algebra of the appropriate pseudo-orthogonal group.) Clifford multiplication by the volume form relates a form to its Hodge dual. In forming Clifford products it is convenient to denote the volume form by z , that is

$$z = *1. \tag{4}$$

We then have

$$*\phi = \phi^\xi z$$

where ξ is the Clifford algebra involution that reverses the order of products and leaves the 1-form generators fixed. Since Hodge duality preserves the space of 2-forms, squaring to minus the identity, the space of complex 2-forms can be decomposed into self-dual and anti-self-dual subspaces satisfying $*F = iF$ and $*F = -iF$ respectively. It then follows that these sub-spaces form simple ideals in the Clifford commutator Lie algebra.

Forms of even degree form a sub-algebra of the Clifford algebra. The complexified even sub-algebra has two inequivalent irreducible representations, the spinor representations. Again we will simply juxtapose symbols to denote the Clifford action on a spinor. So if ψ is a spinor and ω any element of the Clifford algebra we write $\omega\psi$ to denote the Clifford action of ω on ψ . The eigenvalues of the volume form label the inequivalent spinor representations. A spinor ψ is even or odd according to whether $iz\psi = \psi$ or $iz\psi = -\psi$. We have a spin-invariant symplectic product, which we will denote by a bracket $(\ , \)$, that is block diagonal on the inequivalent spinor spaces. This product will be chosen such that

$$(u, \omega v) = (\omega^\xi u, v). \tag{5}$$

This spinor product gives an isomorphism with the space of dual spinors. We let $\bar{u}$ denote the dual spinor such that

$$\bar{u}(v) = (u, v).$$

Since tensor products of spinors and their duals are linear transformations on the space of spinors,

$$(u \otimes \bar{v})w = (v, w)u,$$

we may naturally identify such tensors with elements of the Clifford algebra. Under Clifford multiplication by an arbitrary form ϕ we have

$$\begin{aligned}\phi(u \otimes \bar{v}) &= \phi u \otimes \bar{v} \\ (u \otimes \bar{v})\phi &= u \otimes \overline{\phi^\xi v}.\end{aligned}$$

Under the involution ξ we have

$$(u \otimes \bar{v})^\xi = -v \otimes \bar{u}. \quad (6)$$

The parity of the spinors determines that of their tensor product. For example, if u and v lie in the same spinor space then $u \otimes \bar{v}$ is an even form, as the spinor product is zero on spinors of different parity. Equation (6) shows that the symmetry properties of the tensor product determine the eigenvalue of ξ . For example, the symmetric combination $u \otimes \bar{v} + v \otimes \bar{u}$ is necessarily odd under ξ . So if both spinors lie in the same spinor space this combination is then a 2-form. Moreover, this 2-form will be self-dual if both u and v are even. We may expand a tensor product of spinors into p -form components as

$$u \otimes \bar{v} = \frac{1}{4}(v, u) + \frac{1}{4}(v, e_a u)e^a - \frac{1}{8}(v, e_{ab} u)e^{ab} + \frac{1}{4}(v, e_a z u)e^a z - \frac{1}{4}(v, z u)z. \quad (7)$$

Here we use the abbreviated notation e^{ab} to denote $e^a \wedge e^b$.

Since, in the Lorentzian case, the irreducible representations of the complexified Clifford algebra are the complexifications of those of the real Clifford algebra, we may choose the conjugate linear charge conjugation operator such that it is related to complex conjugation by

$$(u, v)^* = (u^c, v^c)$$

and

$$(u^c)^c = u.$$

In general we follow the conventions of [14].

III. Algebraically-Special Space-times.

In this section we summarise the Petrov classification of the curvature tensor. In particular, we will state the condition that a space-time be algebraically-special in a form convenient to us.

In a four-dimensional Lorentzian space-time the Hodge dual map squares to minus one when acting on 2-forms. With the Hodge dual as complex structure the space of 2-forms may be regarded as a three-dimensional complex space. The Lorentzian metric induces a metric on the space of 2-forms. By using the complex structure of Hodge duality this metric defines a complex Euclidean structure on the space of 2-forms. The curvature tensor can be thought of as a map on the space of 2-forms in such a way that, in an Einstein space, it commutes with Hodge duality and may thus be regarded as a complex linear map. It is also self-adjoint with respect to the complex Euclidean structure. The Petrov classification scheme classifies the Jordan canonical forms of the curvature tensor. Details can be found in [15] and [16].

The metric tensor induces a metric on the space of p -forms. If ϕ and ψ are p -forms then their scalar product $\phi \cdot \psi$ is defined by

$$\phi \wedge * \psi = \phi \cdot \psi * 1,$$

which becomes

$$\phi \cdot \psi = \frac{1}{2} X_a \lrcorner X_b \lrcorner \phi X^a \lrcorner X^b \lrcorner \psi$$

when ϕ and ψ are 2-forms. We may use this metric to regard the tensor product of two 2-forms as an endomorphism on the space of 2-forms;

$$(\phi \otimes \psi)(F) = (\psi \cdot F)\phi.$$

Clearly, symmetric tensor products correspond to self-adjoint operators, and thus more generally so do tensors with ‘pairwise interchange symmetry’. Those tensors that are double-self-dual correspond to endomorphisms on the space of self-dual (or anti-self-dual) 2-forms.

The curvature tensor may be regarded as an endomorphism on 2-forms by using the metric to relate it to a totally covariant tensor,

$$R = 2R^{ab} \otimes e_{ba}$$

where $\{R^{ab}\}$ are the curvature 2-forms. The double-self-dual part of the curvature tensor R^+ is related to the conformal tensor C and the curvature scalar $\mathcal{R}$ by

$$\begin{aligned} R^+ &= C - \frac{1}{6} \mathcal{R} e_{ab} \otimes e^{ab} \\ &= C - \frac{1}{3} \mathcal{R} I \end{aligned}$$

where I is the identity map on 2-forms. The conformal tensor C can be expressed in terms of the conformal 2-forms as

$$C = 2C^{ab} \otimes e_{ba}.$$

Thus acting on an arbitrary 2-form ϕ we have

$$C\phi = 2X_a \lrcorner X_b \lrcorner \phi C^{ab}. \quad (8)$$

The Petrov type of a space-time is determined by the number of eigenvectors and eigenvalues of the conformal tensor when acting in this way on the space of self-dual 2-forms. An algebraically-general conformal tensor has three linearly-independent eigenvectors with distinct eigenvalues, all other cases being classed as algebraically-special.

Any self-dual 2-form ϕ that is null is also decomposable, and hence has one independent real null eigenvector K , $K \lrcorner \phi = 0$. The principal null directions of the conformal tensor correspond to null self-dual 2-forms that satisfy $\phi \wedge C\phi = 0$. (See [15].) This is clearly satisfied by any null eigenform of C . Such eigenforms correspond to repeated principal null directions. An algebraically-special space-time may be characterised as one admitting a null eigenform of the conformal tensor. The only space-times that admit two independent null eigenforms are Petrov type D . In this case the two independent null eigenforms have the same eigenvalue.

IV. Conformal Killing-Yano Tensors.

Killing tensors were introduced as tensors which obey generalisations of Killing's equation, with conformal Killing tensors obeying analogues of the vector conformal Killing equation. One generalisation was to replace the vector with a totally symmetric tensor, whilst Yano [17] extended Killing's equation to a totally anti-symmetric tensor. We shall (as is now common) refer to the latter as Killing-Yano tensors, reserving the term Killing tensor for a totally symmetric tensor.

Killing's equation expresses the condition that a vector field generate an isometry in terms of the symmetrised covariant derivative of the vector field. The Killing tensor and Killing-Yano tensor equations are also usually expressed in terms of the symmetrised covariant derivative. However, the efficiency of the exterior calculus enables the Killing-Yano equation to be written much more succinctly in terms of the anti-symmetrised covariant derivative. If K is any vector field then the covariant derivative ∇K^b can be decomposed into symmetric and skew parts. The symmetric part can be further decomposed into a trace-free part and the trace. The symmetrised covariant derivative is related to the Lie derivative of the metric tensor. In four dimensions we have

$$\nabla_X K^b = \frac{1}{2} X \lrcorner dK^b - \frac{1}{4} X^b d^* K^b + \frac{1}{2} \left\{ \mathcal{L}_K g - \frac{1}{4} \text{Tr} (\mathcal{L}_K g) g \right\} (X)$$

where $\mathcal{L}_K$ denotes the Lie derivative and Tr the trace. (As is usual, we use the metric

tensor to regard any degree two tensor as a linear map on vector fields.) So the conformal-Killing equation can be written (in four dimensions) as

$$\nabla_X K^b = \frac{1}{2} X \lrcorner dK^b - \frac{1}{4} X^b d^* K^b \quad \forall X. \quad (9)$$

If in addition $d^* K^b = 0$ then K is a Killing vector. Notice that (9) implies that

$$e^a \wedge \nabla_{X_a} K^b = \frac{1}{2} e^a \wedge X_a \lrcorner dK^b = dK^b.$$

Since $e^a \wedge \nabla_{X_a} = d$ the coefficient of $X \lrcorner dK^b$ is just such that we cannot conclude that $dK^b = 0$. Similarly (9) implies that

$$X^a \lrcorner \nabla_{X_a} K^b = -\frac{1}{4} X_a \lrcorner (e^a d^* K^b) = -d^* K^b$$

in four dimensions. Thus the coefficient of $X^b d^* K^b$ is just such that we cannot conclude from (9) that $d^* K^b = 0$. This observation suggests how the conformal Killing equation (9) can be generalised to forms of higher degree: the covariant derivative is related to the exterior derivative and the co-derivative with the coefficients chosen such that we do not automatically have the form closed or co-closed. If ω is a 2-form then this generalisation gives the equation

$$3\nabla_X \omega = X \lrcorner d\omega - X^b \wedge d^* \omega \quad \forall X. \quad (10)$$

From this we can use $\nabla_X \omega(Y, Z) = \frac{1}{2} Z \lrcorner Y \lrcorner \nabla_X \omega$ to show that ω must also satisfy

$$\begin{aligned} \nabla_Y \omega(X, Z) + \nabla_Z \omega(X, Y) &= \frac{1}{3} d^* \omega(X) g(Y, Z) \\ &\quad - \frac{1}{6} d^* \omega(Y) g(Z, X) - \frac{1}{6} d^* \omega(Z) g(X, Y) \quad \forall X, Y, Z, \end{aligned} \quad (11)$$

which is Tachibana's conformal generalisation of Yano's Killing equation [2]. Now taking $X = X_a$ and $Y = X_b$ in equation (11), and multiplying both sides by e^{ab} , we recover equation (10). Hence (10) and (11) are equivalent and we may adopt (10) as the conformal Killing-Yano (CKY) equation.

The CKY equation is invariant under Hodge duality. Equation (1) shows that

$$\begin{aligned} *(X^b \wedge d^* \omega) &= -X \lrcorner * d^* \omega \\ &= -X \lrcorner ** d^* \omega \quad \text{by (3)} \\ &= -X \lrcorner d^* \omega \quad \text{by (2)}. \end{aligned}$$

Equivalently

$$*(X \lrcorner d\omega) = (X^b \wedge *d\omega).$$

Since $\nabla_X * = * \nabla_X$ it follows that $*\omega$ is a CKY tensor if ω is. Thus any solution to the CKY equation can be decomposed into self-dual and anti-self-dual CKY tensors.

The CKY equation also often appears in yet another guise. Elements of the Clifford algebra are naturally identified with tensors on the space of spinors. More generally, tensor products of exterior forms may be regarded as higher degree tensors on the spinor spaces, and so any equation for an exterior form can also be written in spinor notation. The CKY equation can be written as

$$\Omega = 0$$

where

$$\Omega = \Omega_a \otimes e^a$$

with

$$\Omega_a = 3\nabla_{X_a} \omega - X_a \lrcorner d\omega + e_a \wedge d^* \omega.$$

The tensor Ω can be thought of as a tensor acting on three even spinors u, v and w , and one odd spinor α by

$$\Omega(u, v, w, \alpha) = (u, \Omega_a v)(\alpha, e^a w).$$

Since the spin-invariant product satisfies (5) we have

$$\begin{aligned} (u, \Omega_a v) &= -(\Omega_a u, v) && \text{for } \Omega_a \text{ 2-forms,} \\ &= (v, \Omega_a u) && \text{since the product is symplectic.} \end{aligned}$$

Thus Ω is automatically symmetric in the first two spinors. It will be totally symmetric in the three even spinors if it is symmetric under interchange of u and w say. Since the space of even spinors is two-dimensional and the product is symplectic we have the identity

$$(u, v)w + (v, w)u + (w, u)v = 0 \quad \text{for all even } u, v, w.$$

Thus

$$\begin{aligned} \Omega(u, v, w, \alpha) &= (u, \Omega_a v)(\alpha, e^a w) \\ &= -(\alpha, e^a \{(\Omega_a v, w)u + (w, u)\Omega_a v\}) \\ &= (v, \Omega_a w)(\alpha, e^a u) - (w, u)(\alpha, e^a \Omega_a v) \\ &= \Omega(v, w, u, \alpha) - (w, u)(\alpha, e^a \Omega_a v) \\ &= \Omega(w, v, u, \alpha) - (w, u)(\alpha, e^a \Omega_a v). \end{aligned}$$

Now

$$e^a \Omega_a = e^a \wedge \Omega_a + X^a \lrcorner \Omega_a,$$

and the Ω_a are such that $e^a \wedge \Omega_a = 0$ and $X^a \lrcorner \Omega_a = 0$. Thus Ω is totally symmetric on the three even spinors. If ω is self-dual then $\Omega(u, v, w, \alpha)$ will be zero unless the first three

spinors are even and the last is odd. Thus the CKY equation for a self-dual 2-form can be regarded as a Spin-irreducible tensor-spinor equation. This spinor equation, equivalent to the CKY equation, was introduced in its own right in [18], and is now usually known as the Killing spinor equation [13]. Because Killing spinors were introduced separately their correspondence with CKY tensors has not always been made clear. There is potential for confusion in that tensors corresponding to Killing spinors have also been called Penrose-Floyd tensors [19].

In the following section we will consider equations for 2-forms related to shear-free congruences. The CKY equation can be regarded as a special case of the shear-free equation. Integrability conditions for the CKY equation then follow as special cases of those for the shear-free equation which will be derived in the following section.

V. Shear-free Equations.

A congruence of curves may be specified by a vector field, the tangent field. The congruence is shear-free if the tangent field generates conformal transformations on its conjugate space (the space of vectors to which it is orthogonal). Thus the shear-free condition is a generalisation of the conformal-Killing condition. Since the shear-free condition is conformally invariant, the various shear-free equations that will be given all have a conformal covariance. If a vector field generates conformal transformations on its conjugate then so does any vector field proportional to it. Since a reparametrisation of the congruence corresponds to a scaling of the tangent field the shear-free condition is a reparametrisation-invariant property of the congruence. The condition that a congruence be shear-free can be formulated as a ‘gauged’ conformal Killing equation, where the connection terms ensure covariance under a scaling of the vector field. Thus a vector field K corresponds to a shear-free congruence if it satisfies the equation [20]

$$\widehat{\nabla}_X K^b = \frac{1}{2} X \lrcorner \hat{d} K^b - \frac{1}{4} X^b \hat{d}^* K^b \quad \forall X. \quad (12)$$

Here $\widehat{\nabla}$ is a scaling-covariant derivative,

$$\widehat{\nabla}_X K^b = \nabla_X K^b + 2qA(X)K^b$$

for some real 1-form A . (The factor of 2 and the constant q will be convenient later.) The gauged exterior derivative $\hat{d}$ and co-derivative $\hat{d}^*$ are related to $\widehat{\nabla}$ by

$$\begin{aligned} \hat{d} &= e^a \wedge \widehat{\nabla}_{X_a} \\ \hat{d}^* &= -X^a \lrcorner \widehat{\nabla}_{X_a} . \end{aligned}$$

(Equation (12) has the numerical coefficients chosen for four dimensions, although it is easily generalised to arbitrary dimensions.) Throughout we shall write various shear-free

equations in terms of a ‘gauged’ covariant derivative. However, it must be remembered that the form A , playing the role of connection, is not some given background field, but depends upon the vector field K . When K is non-null then A can be expressed in terms of K , whereas in the null case only certain components can be expressed in terms of K [20].

If K is null (and real) then it may be related to an even spinor u by

$$K^b = (iu^c, e^a u) e_a.$$

Clearly the correspondence between K and u is not one-to-one, there being a $U(1)$ freedom in the choice of u . The shear-free condition (12) for K is then equivalent to the following equation for u :

$$\widehat{\nabla}_X u - \frac{1}{4} X^b \widehat{D} u = 0 \quad \forall X \quad (13)$$

where

$$\widehat{\nabla}_X u = \nabla_X u + q\mathcal{A}(X)u \quad (14)$$

where $\mathcal{A}$ is a complex 1-form whose real part is A , and $\widehat{D}$ is the Dirac operator

$$\widehat{D} = e^a \widehat{\nabla}_{X_a}.$$

The shear-free spinor equation (13) is a $\mathbb{C}^*$ -covariant twistor equation, where $\mathbb{C}^*$ is the group of non-zero complex numbers. The $U(1)$ part of the covariance stems from the projective relationship between u and K , whilst the scaling part of the covariance is related to the reparametrisation-invariance of the shear-free condition. In the same way that we showed the spinorial correspondence of the CKY equation in the previous section, we may regard (13) as an equation for a spin tensor acting symmetrically on two even spinors and one odd spinor. Written thus the shear-free spinor equation was obtained by Sommers [21].

A (real) null vector field K may be put into correspondence with a self-dual decomposable 2-form ϕ by the relation

$$K^b \phi = 0.$$

A given K only determines ϕ up to a complex scaling. The shear-free condition for K gives rise to an equation for ϕ . In fact Robinson’s theorem [3] shows that ϕ is proportional to a closed (and hence, since it is self-dual, co-closed) 2-form; that is, a Maxwell solution. It will be convenient for us to state the shear-free condition for ϕ differently. In terms of the even spinor u representing K we may choose

$$\phi = u \otimes \bar{u}. \quad (15)$$

The shear-free condition for u then translates to an equation for ϕ . It is simplest to obtain the corresponding equation for ϕ written in terms of a Clifford commutator, [,]. If we write ϕ as in (15) then we can show that u satisfies (13) if and only if

$$\widehat{\nabla}_X \phi - \frac{1}{4} [X^b e^a, \widehat{\nabla}_{X_a} \phi] = 0 \quad (16)$$

where

$$\widehat{\nabla}_X \phi = \nabla_X \phi + 2q\mathcal{A}(X)\phi. \quad (17)$$

Since the Clifford commutator term can be written as

$$\frac{1}{2}[X^b e^a, \widehat{\nabla}_{X_a} \phi] = X \lrcorner \hat{d}\phi - X^b \wedge \hat{d}^* \phi - \widehat{\nabla}_X \phi$$

we see that the shear-free equation (13) is equivalent to the gauged conformal Killing-Yano equation

$$3\widehat{\nabla}_X \phi = X \lrcorner \hat{d}\phi - X^b \wedge \hat{d}^* \phi. \quad (16')$$

(We reiterate that the form $\mathcal{A}$ depends upon the 2-form ϕ : in particular, should there be two shear-free 2-forms then, in general, the ‘gauge terms’ occurring in each equation will be different.) Thus as an alternative to Robinson’s theorem we can state that a decomposable self-dual 2-form corresponds to a shear-free null congruence if and only if it satisfies the $\mathbb{C}^*$ -gauged conformal Killing-Yano equation. One can show that null solutions to (16') can be scaled to produce Maxwell fields, and vice versa, and so indeed this statement is equivalent to Robinson’s theorem. Previously Dietz and Rüdiger [4] investigated a generalisation of Robinson’s theorem. They considered non-decomposable 2-forms corresponding to two independent null congruences. They showed that both these congruences are shear-free if and only if the self-dual 2-form satisfies an equation equivalent to (16'). Although they showed that their equation was a generalisation of the CKY equation they did not interpret the extra terms as gauge terms. Moreover, they only considered non-decomposable solutions to the equation. In fact it is rather nice to have a single equation for a 2-form that characterises any eigenvectors as being shear-free, whether there be one or two independent real eigenvectors.

The shear-free equations can be differentiated to obtain integrability conditions relating second derivatives to curvature terms. We shall make use of these later on. Firstly we consider the shear-free spinor equation. Differentiating (13) introduces the curvature operator $\hat{R}(X, Y)$ of $\widehat{\nabla}$. Since $\widehat{\nabla}$ is related to ∇ by (14) the curvature operators are related by

$$\hat{R}(X, Y)u = R(X, Y)u - qX \lrcorner Y \lrcorner \mathcal{F}u \quad (18)$$

where $\mathcal{F}$ is the $\mathbb{C}^*$ curvature,

$$\mathcal{F} = d\mathcal{A}.$$

Since

$$R(X, Y)u = \frac{1}{2}e^a(X)e^b(Y)R_{ab}u \quad (19)$$

where R_{ab} are the curvature 2-forms, equation (13) has the integrability condition

$$R_{ab}u + 2qX_b \lrcorner X_a \lrcorner \mathcal{F}u - \frac{1}{2}(e_b \widehat{\nabla}_{X_a} - e_a \widehat{\nabla}_{X_b})\hat{D}u = 0. \quad (20)$$

Multiplying by e_a gives

$$P_b u - 2qX_b \lrcorner \mathcal{F}u + \hat{\nabla}_{X_b} \hat{D}u + \frac{1}{2}e_b \hat{D}^2 u = 0 \quad (21)$$

where P_b are the Ricci 1-forms [14]. Multiplying this by e^b produces

$$\hat{D}^2 u = \frac{4q}{3} \mathcal{F}u - \frac{1}{3} \mathcal{R}u \quad (22)$$

where $\mathcal{R}$ is the curvature scalar. A Laplacian on spinors is given by the trace of the Hessian,

$$\hat{\nabla}^2 = \hat{\nabla}_{X^a} \hat{\nabla}_{X_a} - \hat{\nabla}_{\nabla_{X_a} X^a}.$$

This is related to the square of the Dirac operator by curvature terms:

$$\hat{D}^2 u = \hat{\nabla}^2 u + \frac{1}{2} e^{ab} \hat{R}(X_a, X_b)u. \quad (23)$$

By (18) and (19) we have

$$e^{ab} \hat{R}(X_a, X_b)u = -\frac{1}{2} \mathcal{R}u + 2q \mathcal{F}u$$

and so (22) can be written as

$$\hat{\nabla}^2 u = \frac{q}{3} \mathcal{F}u - \frac{1}{12} \mathcal{R}u. \quad (22')$$

Since the conformal 2-forms C_{ab} are given by

$$C_{ab} = R_{ab} - \frac{1}{2}(P_a \wedge e_b - P_b \wedge e_a) + \frac{1}{6} \mathcal{R}e_a \wedge e_b$$

we may use (20), (21) and (22) to obtain the integrability condition

$$C_{ab}u + q\Gamma_{ab}u = 0 \quad (24)$$

where

$$\Gamma_{ab} = \frac{1}{6} e_{ba} \mathcal{F} + \frac{1}{2} \mathcal{F}e_{ba}. \quad (25)$$

Now we look at the integrability conditions for the decomposable self-dual 2-form describing the shear-free congruence. If ϕ is related to u by (15) then we may use the integrability

condition (24) of (13) to obtain an analogous integrability condition of (16'). If we define $C\phi$ by (8) then from (15) and (7)

$$\begin{aligned} C\phi &= -\frac{1}{2}(u, e_{ab}u)C^{ba} \\ &= -\frac{1}{2}(u, C_{ab}u)e^{ba} \end{aligned}$$

by the ‘pairwise symmetry’ of the conformal tensor. Now we may use (24) to obtain

$$\begin{aligned} C\phi &= \frac{q}{2}(u, \Gamma_{ab}u)e^{ba} \\ &= -\frac{q}{4}(u, (\Gamma_{ab} - \Gamma_{ab}^\xi)u)e^{ab}. \\ &= -\frac{q}{12}(u, [\mathcal{F}, e_{ba}]u)e^{ab} \end{aligned}$$

from (25),

$$= -\frac{q}{12}(u, e_{ab}u)[\mathcal{F}, e^{ab}]$$

due to the ‘pairwise anti-symmetry’ of $X_p \lrcorner X_q \lrcorner [\mathcal{F}, e_{ba}]$,

$$= \frac{2q}{3}[\mathcal{F}, \phi] \tag{26}$$

from (7).

In analogy with (23) the trace of the Hessian is related to the ‘gauged’ Laplace-Beltrami operator $\hat{\Delta}$ by

$$\begin{aligned} \hat{\Delta}\phi &= \hat{\nabla}^2\phi + e^b \wedge X^a \lrcorner \hat{R}(X_b, X_a)\phi \\ &= \hat{\nabla}^2\phi + \frac{1}{4}[e^{ba}, \hat{R}(X_b, X_a)]\phi \end{aligned}$$

where

$$\hat{\Delta}\phi = -(\hat{d}\hat{d}^* + \hat{d}^*\hat{d})\phi.$$

Since $\hat{\nabla}$ is related to ∇ by (17) the curvatures are related by

$$\hat{R}(X, Y)\phi = R(X, Y)\phi - 2qX \lrcorner Y \lrcorner \mathcal{F}\phi$$

and hence

$$\begin{aligned} [e^{ba}, \hat{R}(X_b, X_a)]\phi &= [e^{ba}, R(X_b, X_a)]\phi + 4q[\mathcal{F}, \phi] \\ &= -2C\phi - \frac{4}{3}\mathcal{R}\phi + 4q[\mathcal{F}, \phi]. \end{aligned}$$

Thus $\hat{\Delta}$ is related to $\hat{\nabla}^2$ by

$$\hat{\Delta}\phi = \hat{\nabla}^2\phi - \frac{1}{2}C\phi - \frac{1}{3}\mathcal{R}\phi + q[\mathcal{F}, \phi]. \tag{27}$$

By differentiating (16') we obtain

$$3\hat{\nabla}^2\phi = \hat{\Delta}\phi,$$

and so (26) gives the integrability condition

$$\hat{\nabla}^2\phi = \frac{q}{3}[\mathcal{F}, \phi] - \frac{1}{6}\mathcal{R}\phi \quad (28)$$

or

$$\hat{\Delta}\phi + \frac{1}{2}\mathcal{R}\phi = q[\mathcal{F}, \phi], \quad (28')$$

where the left-hand side is the conformally covariant (gauged) Laplace-Beltrami operator.

The integrability condition (26) shows that $\phi \wedge C\phi = 0$ and thus the null ϕ corresponds to a principal direction. In a Ricci-flat space-time the Goldberg-Sachs theorem makes the stronger statement that a shear-free null congruence must correspond to a *repeated* principal null direction, and conversely any repeated principal null direction must correspond to a shear-free congruence [22, 23]. More generally necessary and sufficient conditions for a shear-free congruence to correspond to a repeated principal null direction are given by the generalised Goldberg-Sachs theorem [24]. So in the generalised Goldberg-Sachs class of space-times ϕ must be an eigenform of the conformal tensor. Thus from (26) the commutator of $\mathcal{F}$ and ϕ must be proportional to ϕ , and so from (15) we see that the spinor u must be an eigenvector of $\mathcal{F}$. Thus

$$q\mathcal{F}u = 3\mu u, \quad (29)$$

for some eigenfunction μ , and

$$q[\mathcal{F}, \phi] = 6\mu\phi. \quad (30)$$

The integrability condition (22) for the spinor becomes

$$\hat{D}^2u = 4\mu u - \frac{1}{3}\mathcal{R}u \quad (31a)$$

$$\hat{\nabla}^2u = \mu u - \frac{1}{12}\mathcal{R}u, \quad (31b)$$

whilst (28) becomes

$$\hat{\nabla}^2\phi = 2\mu\phi - \frac{1}{6}\mathcal{R}\phi \quad (32a)$$

$$\hat{\Delta}\phi + \frac{1}{2}\mathcal{R}\phi = 6\mu\phi, \quad (32b)$$

and (26) becomes

$$C\phi = 4\mu\phi. \quad (33)$$

We can obtain integrability conditions for the CKY equation using the results that we have for shear-free equations. In the case of a null self-dual CKY tensor we simply have the conditions above where the ‘gauge’ terms are zero. In this case the corresponding shear-free spinor equation reduces to the twistor equation. From (26) we see that the CKY tensor is a null eigenvector of C with eigenvalue zero. In fact (24) shows that any self-dual 2-form ϕ that has u as eigenspinor satisfies $C\phi = 0$. Therefore C must have two linearly-independent eigenvectors, and so the space-time must be type N or conformally-flat.

If ω is a non-null self-dual 2-form then we may write ω in terms of a pair of even spinors u_1 and u_2 as

$$\omega = \frac{1}{2}(u_1 \otimes \bar{u}_2 + u_2 \otimes \bar{u}_1). \quad (34)$$

In the same way that we showed that the shear-free spinor equation (13) led to the ‘gauged’ conformal Killing-Yano equation (16) we may show that the non-null ω satisfies the CKY equation if and only if the spinors $\{u_i\}$ both satisfy a shear-free equation, with the spinors having opposite $\mathbb{C}^*$ ‘charges’. (Dietz and Rüdiger showed that the 2-form ω satisfies a gauged CKY equation if and only if the spinors $\{u_i\}$ satisfy the shear-free equation [4]. In general the $\mathbb{C}^*$ ‘charge’ of ω is the sum of those of u_1 and u_2 . So we have here just a special case when ω is a CKY tensor.)

We may use the integrability conditions for the spinor equations to obtain integrability conditions for ω , just as we did in the shear-free case. We have

$$\begin{aligned} C\omega &= -\frac{1}{2}(u_2, e_{ab}u_1)C^{ba} \\ &= -\frac{1}{2}(u_2, C_{ab}u_1)e^{ba} \\ &= -\frac{1}{2}(u_1, C_{ab}u_2)e^{ba}. \end{aligned}$$

The integrability condition (24) for u_1 gives

$$C\omega = \frac{q_1}{2}(u_2, \Gamma_{ab}u_1)e^{ba}, \quad (35)$$

whilst that for u_2 gives

$$\begin{aligned} C\omega &= \frac{q_2}{2}(u_1, \Gamma_{ab}u_2)e^{ba} \\ &= \frac{q_1}{2}(u_2, \Gamma_{ab}^\xi u_1)e^{ba} \end{aligned}$$

since $q_2 = -q_1$.

Subtracting this from (35) shows that

$$(u_2, (\Gamma_{ab} - \Gamma_{ab}^\xi)u_1)e^{ba} = 0.$$

Repeating the steps that lead to (26) then shows that we must have

$$[\mathcal{F}, \omega] = 0. \quad (36)$$

From this we can conclude that the self-dual part of $\mathcal{F}$ is proportional to ω . Since u_1 and u_2 are eigenvectors of ω , and hence $\mathcal{F}$, there must be some function μ such that

$$q_i \mathcal{F} u_i = 3\mu u_i \quad (\text{no sum.}) \quad (29')$$

Both of the $\{u_i\}$ satisfy the integrability conditions (31a) for the appropriate covariant derivative. From each of the spinors we can make a null shear-free 2-form:

$$\phi_i = u_i \otimes \bar{u}_i \quad (\text{no sum,})$$

each satisfying the integrability condition (33). Thus each of the ϕ_i corresponds to a repeated principal null direction. Thus for a non-null CKY tensor to exist we must have Petrov type D (or conformally-flat), within the generalised Goldberg-Sachs class of space-times. Each of ϕ_1 and ϕ_2 satisfies the integrability conditions (32a).

We now return to the integrability condition (35) for ω . If we insert (25) into (35) and use (29') then we have

$$C\omega = -8\mu\omega. \quad (37)$$

Thus ϕ_1 , ϕ_2 and ω form an eigenbasis of self-dual 2-forms under the action of the conformal tensor. The CKY tensor ω also satisfies (32a) where the 'gauge terms' are absent from the Laplace-Beltrami operator.

We have seen that for a non-null CKY tensor to exist the space-time is necessarily Petrov type D . Suppose now that we have Petrov type D with ϕ_1 and ϕ_2 null shear-free 2-forms corresponding to repeated principal null directions. We cannot, in general, assume any relation between the gauge terms in the two different shear-free equations. We let ω be defined by (34) such that $\{\phi_1, \phi_2, \omega\}$ is an eigenbasis for C . Since each ϕ_i is an eigenvector of C (29) becomes

$$q_i \mathcal{F}_i u_i = 3\mu_i u_i \quad (\text{no sum,}) \quad (29'')$$

where now we cannot assume that $\mathcal{F}_1$ and $\mathcal{F}_2$ are related. It then follows that each ϕ_i is an eigenvector of C with eigenvalue $4\mu_i$, and since the eigenvalues are assumed equal we

can conclude that $\mu_1 = \mu_2$. Proceeding as we did in (35) we can express $C\omega$ in terms of the integrability conditions for either spinor:

$$\begin{aligned} C\omega &= \left(\frac{\mu}{4}(u_2, e_{ba}u_1) - \frac{1}{4}(q_1\mathcal{F}_1u_2, e_{ba}u_1) \right) e^{ba} \\ &= \left(\frac{\mu}{4}(u_2, e_{ba}u_1) - \frac{1}{4}(q_2\mathcal{F}_2u_1, e_{ba}u_2) \right) e^{ba}. \end{aligned}$$

Since ω has been assumed an eigenvector of C we must have u_2 an eigenspinor of $\mathcal{F}_1$. By taking the product with u_1 , which is also an eigenspinor, we see that the eigenvalues of the two spinors must be opposite. We can draw similar conclusions for $\mathcal{F}_2$ and have

$$\begin{aligned} q_1\mathcal{F}_1u_1 &= 3\mu u_1 \\ q_1\mathcal{F}_1u_2 &= -3\mu u_2 \\ q_2\mathcal{F}_2u_1 &= -3\mu u_1 \\ q_2\mathcal{F}_2u_2 &= 3\mu u_2. \end{aligned}$$

It follows that the self-dual part of $q_1\mathcal{F}_1 + q_2\mathcal{F}_2$ is zero. That is, the self-dual part of the curvature of the gauge term entering into the equation for ω is zero (as was shown by Dietz and Rüdiger [4]). It then follows that ω satisfies (27) with the $\mathbb{C}^*$ -curvature term on the right-hand side zero. Hence ω satisfies (32a), and in addition (30). Thus (32a) is satisfied by any shear-free 2-form whose eigenvectors are aligned with repeated principal directions.

We have seen that in type D space-times the self-dual part of the $\mathbb{C}^*$ -curvature associated with ω must vanish. When the anti-self-dual part is also zero then ω can be scaled to a CKY tensor. Penrose and Walker's result on Killing spinors shows that this can be done in every type D Einstein space [18].

VI. Debye potentials and symmetry operators for source-free Maxwell fields.

Maxwell's equations require that the electromagnetic field 2-form F be closed. This is automatically satisfied if F is exact, $F = dA$ say, where A is the potential 1-form. In source-free regions F is also required to be co-closed, that is $d^*F = 0$. This would be automatically satisfied if there were a co-potential 3-form B such that $F = d^*B$. Suppose that a 2-form Z satisfies

$$\Delta Z = -d\mathcal{G} - d^*\mathcal{W}, \tag{38}$$

where $\mathcal{G}$ and $\mathcal{W}$ are arbitrary forms of degree one and three respectively. Then by noting that this can be rewritten as

$$d^*(\mathcal{W} - dZ) = d(d^*Z - \mathcal{G}) \tag{38'}$$

we see that Z provides us with a 2-form that is both exact and co-exact and hence a Maxwell solution. The 2-form Z is called a Hertz potential [25]. On the face of it (38) doesn't offer a promising approach, solving this equation is no easier than solving the original Maxwell equations. However, in some circumstances, with an appropriate choice of the form for Z , solutions to (38) can be written in terms of solutions to a scalar equation; a Debye potential equation [26, 1 and references therein]. Cohen and Kegeles [1] applied the Debye potential method to the solution of Maxwell's equations in curved space-times. They pointed out that in algebraically-special space-times there exist privileged 2-forms corresponding to the repeated principal null direction. Using the Newman-Penrose formalism they explicitly obtained scalar Debye potentials by aligning the Hertz potential with the repeated principal null direction: the resulting scalar equation being expressed in the adapted null basis.

In this section we show how a Debye potential is obtained by choosing Z to be proportional to a shear-free 2-form in an algebraically-special space-time. As we shall explicitly use the shear-free equation, rather than an adapted basis, the resulting Debye potential equation will be expressed in a basis-independent form. The scheme becomes more powerful when a conformal Killing-Yano tensor exists.

Consider a shear-free 2-form ϕ , satisfying (16'). (We allow ϕ to be either null or non-null, and in the case in which the $\mathbb{C}^*$ -charge is zero then we have a CKY tensor.) Let f be a scalar field with opposite $\mathbb{C}^*$ -charge so that the 2-form $f\phi$ is a $\mathbb{C}^*$ -scalar. Then the Laplacian of $f\phi$ can be expressed as

$$\nabla^2(f\phi) = \hat{\nabla}^2 f\phi + 2\hat{\nabla}_{X_a} f \hat{\nabla}^{X_a} \phi + f \hat{\nabla}^2 \phi$$

where $\hat{\nabla}$ is the $\mathbb{C}^*$ -covariant derivative. We can write this in terms of the the $\mathbb{C}^*$ -gradient as

$$\nabla^2(f\phi) = \hat{\nabla}^2 f\phi + 2\hat{\nabla} \widehat{\text{grad}}_f \phi + f \hat{\nabla}^2 \phi,$$

or

$$\Delta(f\phi) = \hat{\nabla}^2 f\phi + 2\hat{\nabla} \widehat{\text{grad}}_f \phi + f \hat{\nabla}^2 \phi - \frac{1}{2} f C \phi - \frac{1}{3} f \mathcal{R} \phi \quad (39)$$

by (27). Now, from the shear-free equation (16') we have

$$\begin{aligned} 3\hat{\nabla} \widehat{\text{grad}}_f \phi &= \widehat{\text{grad}} f \lrcorner \hat{d}\phi - \hat{d}f \wedge \hat{d}^* \phi \\ &= X^a \lrcorner \left(\hat{\nabla}_{X_a} (f \hat{d}\phi) - f \hat{\nabla}_{X_a} \hat{d}\phi \right) - d(f \hat{d}^* \phi) + f \hat{d} \hat{d}^* \phi \\ &= -d^*(f \hat{d}\phi) + f \hat{d}^* \hat{d}\phi - d(f \hat{d}^* \phi) + f \hat{d} \hat{d}^* \phi \\ &= -d^*(f \hat{d}\phi) - d(f \hat{d}^* \phi) - f \hat{\Delta} \phi. \end{aligned}$$

Inserting this into (39) gives

$$\Delta(f\phi) = \hat{\nabla}^2 f\phi - \frac{2}{3} d^* (f \hat{d}\phi) - \frac{2}{3} d (f \hat{d}^* \phi) + f \left(\hat{\nabla}^2 \phi - \frac{2}{3} \hat{\Delta} \phi \right) - \frac{1}{2} f C \phi - \frac{1}{3} f \mathcal{R} \phi.$$

When ϕ is self-dual with its eigenvectors repeated principal null directions this becomes, by (32a),

$$\Delta(f\phi) = \left(\hat{\nabla}^2 f - \frac{1}{6} \mathcal{R}f \right) \phi - \lambda_\phi f \phi - \frac{2}{3} d^* (f \hat{d}\phi) - \frac{2}{3} d (f \hat{d}^* \phi)$$

where

$$2\mu\phi + \frac{1}{2}C\phi = \lambda_\phi\phi. \quad (40)$$

It is convenient to identify λ_ϕ as an eigenvalue,

$$C\phi = n^2 \lambda_\phi \phi, \quad (40')$$

where n is the number of (real) eigenvectors of ϕ . Thus when ϕ is null $n = 1$ and λ_ϕ is just the eigenvalue of C corresponding to ϕ , whereas in the non-null case $n = 2$ and λ_ϕ is a quarter of the eigenvalue of C . So if the scalar field satisfies the equation

$$\hat{\nabla}^2 f - \frac{1}{6} \mathcal{R}f = \lambda_\phi f \quad (41)$$

then we can relate an exact form to a co-exact one:

$$d \left(d^*(f\phi) - \frac{2}{3} f \hat{d}^* \phi \right) = d^* \left(\frac{2}{3} f \hat{d}\phi - d(f\phi) \right).$$

Thus out of the 2-form ϕ and the scalar f we have a source-free Maxwell solution,

$$F'(f, \phi) = d \left(d^*(f\phi) - \frac{2}{3} f \hat{d}^* \phi \right) \quad (42a)$$

$$= d^* \left(\frac{2}{3} f \hat{d}\phi - d(f\phi) \right). \quad (42b)$$

Notice how with this approach, unlike that of Cohen and Kegeles, we do not need to choose the gauge terms $\mathcal{G}$ and $\mathcal{W}$ by inspection, rather they appear naturally and are given explicitly in term of ϕ . Using (3) and the fact that $\hat{d}^* = *\hat{d}*$, we can see that $F'(f, \phi)$ is anti-self-dual for self-dual ϕ .

To apply this scheme to construct a Maxwell solution in a given space-time it is necessary to solve the scalar equation (41). The shear-free 2-form ϕ enters into this equation, not only through the eigenfunction λ_ϕ , but also through the ‘gauge term’ $\mathcal{A}$. We may expose the gauge terms in the shear-free equation (16') by writing it as

$$Y_X \phi = q[X^\flat \wedge \mathcal{A}, \phi] - 4q\mathcal{A}(X)\phi \quad (43)$$

where we have defined Y_X by

$$Y_X \phi = 3\nabla_X \phi - X \lrcorner d\phi + X^\flat \wedge d^* \phi. \quad (44)$$

When there is only one repeated principal null direction then the self-dual ϕ will be null. To use (43) to determine $\mathcal{A}$ it is convenient to pick some adapted basis. (Such a point comes to us all when we have to actually solve equations.) Let Σ be the maximal totally isotropic subspace such that $X \in \Sigma \Leftrightarrow X^\flat \phi = 0$. Then for X in Σ (43) reduces to

$$Y_X \phi = -2q\mathcal{A}(X)\phi \quad \forall X \in \Sigma.$$

This enables half of the components of $\mathcal{A}$ to be found. To compute the remaining components we may pick a maximal isotropic subspace Σ' to complement Σ . Then let ψ be a null self-dual 2-form such that $X \in \Sigma' \Leftrightarrow X^\flat \psi = 0$. Then the remaining components of $\mathcal{A}$ can be calculated from (43) which becomes

$$Y_X \phi \cdot \psi = -6q\mathcal{A}(X)\phi \cdot \psi \quad \forall X \in \Sigma'.$$

In a type D space-time there are two repeated principal null directions. Thus one can choose either of the two corresponding null shear-free 2-forms as Hertz potentials, or one can choose the non-null 2-form ω that has both repeated principal directions as eigenvectors. Thus there are three possible choices of Hertz potential, as was pointed out by Mustafa and Cohen [27]. They chose to normalise ω to have constant length, whereas we have chosen to scale ω to be CKY and so their equation for a non-null Debye potential will differ from ours by exact gauge terms.

In the non-null case the gauge term is determined directly in terms of ω . In this case (43) gives

$$Y_X \omega \cdot \omega = -4q\mathcal{A}(X)\omega \cdot \omega \quad \forall X.$$

This expression was given by Dietz and Rüdiger [4]. Clearly the scheme simplifies in the case in which we have a CKY tensor, for then the gauge terms are absent from the Debye potential equation.

Equation (42a) enables a Maxwell field to be constructed from a shear-free 2-form and a scalar Debye potential satisfying (41). It turns out that we can conversely take a shear-free 2-form and a Maxwell field and construct a scalar Debye potential. If $f(F, \phi) = F \cdot \phi$ then the $\mathbb{C}^*$ -covariant Laplacian is given by

$$\hat{\nabla}^2 f(F, \phi) = \nabla^2 F \cdot \phi + 2\nabla_{X_a} F \cdot \hat{\nabla}_{X^a} \phi + F \cdot \hat{\nabla}^2 \phi,$$

where the $\mathbb{C}^*$ -charge of $f(F, \phi)$ is the same as that of ϕ . Since ϕ satisfies the shear-free equation (16')

$$\begin{aligned} 3\nabla_{X_a} F \cdot \hat{\nabla}_{X^a} \phi &= \nabla_{X_a} F \cdot (X^a \lrcorner \hat{d}\phi) - \nabla_{X_a} F \cdot (e^a \wedge \hat{d}^* \phi) \\ &= dF \cdot \hat{d}\phi + d^* F \cdot \hat{d}^* \phi = 0 \end{aligned}$$

by Maxwell's equations. We may then use (27) to relate $\nabla^2 F$ to ΔF , which is zero by Maxwell's equations, to obtain

$$\begin{aligned}\hat{\nabla}^2 f(F, \phi) &= \left(\frac{1}{2}CF + \frac{1}{3}\mathcal{R}F \right) \cdot \phi + F \cdot \hat{\nabla}^2 \phi \\ &= F \cdot \left(\frac{1}{2}C\phi + \frac{1}{3}\mathcal{R}\phi + \hat{\nabla}^2 \phi \right)\end{aligned}$$

since C is self-adjoint,

$$= F \cdot \left(\frac{1}{2}C\phi + \frac{1}{6}\mathcal{R}\phi + 2\mu\phi \right)$$

by (32a),

$$= F \cdot (\lambda_\phi \phi + \frac{1}{6}\mathcal{R}\phi).$$

So $f(F, \phi)$ satisfies equation (41): notice, however, that $f(F, \phi)$ has the same 'charge' as ϕ , whereas the construction of a Maxwell field from ϕ requires a Debye potential with opposite charge.

For those space-times that admit a non-null CKY tensor the above gives a method of mapping any Maxwell field to another. As was shown in section 5, when we have a non-null CKY tensor ω we have a pair of null shear-free 2-forms, ϕ_1 and ϕ_2 , having opposite $\mathbb{C}^*$ -charge. We can take one of these, ϕ_1 say, and a given Maxwell field F to construct a Debye potential $f(F, \phi_1)$. This will then have the appropriate charge to combine with ϕ_2 to form a new Maxwell field (also note that $\lambda_{\phi_1} = \lambda_{\phi_2}$). That is, we have a symmetry operator $\mathcal{L}_{\phi_1\phi_2}$, mapping between Maxwell fields, defined by

$$\mathcal{L}_{\phi_1\phi_2} F = F'(F \cdot \phi_1, \phi_2).$$

Interchanging the roles of ϕ_1 and ϕ_2 gives another symmetry operator. However, by using the shear-free equations for ϕ_1 and ϕ_2 it can be shown that when acting on a Maxwell field F , their difference vanishes and so

$$\mathcal{L}_{\phi_1\phi_2} F = \mathcal{L}_{\phi_2\phi_1} F.$$

We can also use a non-null CKY ω directly to make a symmetry operator for Maxwell fields. The scalar $f(F, \omega)$ is a Debye potential satisfying an uncharged equation and hence can be combined again with ω to produce a Maxwell field. Hence

$$\mathcal{L}_{\omega\omega} F = F'(F \cdot \omega, \omega)$$

is another symmetry operator. Since self-dual and anti-self-dual 2-forms are mutually orthogonal, these symmetry operators map only the self-dual part of a Maxwell field to an

anti-self-dual Maxwell field. To see the relationship between these symmetry operators, we will first recast them in terms of a higher order generalisation of a CKY 2-form.

The CKY tensor ω enters quadratically in the symmetry operator $\mathcal{L}_{\omega\omega}$. By taking the tensor product of ω with itself we obtain a tensor, quadratic in ω , which we may regard as an endomorphism on the space of 2-forms, as we did in section 3. In this way we may write $\mathcal{L}_{\omega\omega}$ in terms of a degree-four tensor constructed from ω . Let P^+ be the operator that projects out the self-dual part of any 2-form. Then clearly P^+ commutes with the Hodge dual. Since it is self-adjoint with respect to the metric on 2-forms it corresponds to a pairwise-symmetric tensor. Then out of the self-dual ω we construct the tensor K :

$$K = \omega \otimes \omega - \frac{1}{3}(\omega \cdot \omega)P^+. \quad (45)$$

In section 4 we showed how ω corresponded to a spin-tensor. In the same way one can show that K corresponds to a totally symmetric spin tensor. The symmetry operator constructed from ω can be written in terms of K as

$$\mathcal{L}_{\omega\omega}F = d \left(d^*(KF) - \frac{2}{5}D^*K(F) \right) \quad (46a)$$

$$= d^* \left(\frac{2}{5}DK(F) - d(KF) \right) \quad (46b)$$

where the ‘exterior derivative’ D is defined by

$$DK(G) = d(KG) - e^a \wedge K(\nabla_{X_a}G)$$

for G an arbitrary 2-form. The ‘co-derivative’ D^* is defined analogously.

The only non-zero inner products between a CKY ω and its associated oppositely charged null shear-free 2-forms ϕ_1 and ϕ_2 are related by

$$\phi_1 \cdot \phi_2 = -2\omega \cdot \omega.$$

After calculating the action of K on the self-dual basis $\{\omega, \phi_1, \phi_2\}$ we can use this to see that an alternative expression for K is

$$K = \frac{1}{2}\phi_1 \otimes \phi_2 + \frac{1}{2}\phi_2 \otimes \phi_1 - \frac{1}{3}\phi_1 \cdot \phi_2 P^+.$$

Then using the shear-free equations for ϕ_1 and ϕ_2 we can rewrite the right hand side of (46a) to show that

$$\begin{aligned} \mathcal{L}_{\omega\omega}F &= \frac{1}{2}\mathcal{L}_{\phi_1\phi_2}F + \frac{1}{2}\mathcal{L}_{\phi_2\phi_1}F \\ &= \mathcal{L}_{\phi_1\phi_2}F = \mathcal{L}_{\phi_2\phi_1}F, \end{aligned}$$

since we have already seen that the two terms on the right hand side are equal when acting on Maxwell fields. Hence, as was pointed out by Torres del Castillo [11] who wrote down these operators using the two-component spinor formalism, the various Debye schemes give rise only one symmetry operator.

The CKY equation for ω can be used to obtain an analogous equation for K . The analogy is closest if we write the CKY equation in terms of the Clifford commutator as in (16). For any 2-form ϕ let L_ϕ be the operator that maps any 2-form to the Clifford commutator: $L_\phi(\psi) = [\phi, \psi]$. Then the CKY equation can be written as

$$\nabla_X \omega - \frac{1}{4} L_{X^\flat \wedge e^a} \nabla_{X_a} \omega = 0,$$

whilst K satisfies the equation

$$\nabla_X K - \frac{1}{6} [L_{X^\flat \wedge e^a}, \nabla_{X_a} K] = 0. \quad (47)$$

Here the bracket denotes the commutator of the operators. One can show that equation (47) corresponds to a spin-irreducible spin-tensor equation, as we showed is the case for the CKY equation. Just as the CKY equation corresponds to the Killing spinor equation, equation (47) corresponds to the ‘4-index Killing spinor’ equation. Kalnins, McLenaghan and Williams obtained a symmetry operator for Maxwell fields from a 4-index Killing spinor [12]. They then obtained a corresponding tensor equation which they observed was analogous to the CKY equation as written by Tachibana [2] (equation (11)).

VII. Debye potentials and symmetry operators for massless Dirac fields.

In the previous section we showed how we could associate a Debye potential with a shear-free 2-form, enabling the source-free Maxwell equations to be solved in terms of solutions to a scalar equation. In the more special case in which there existed a CKY tensor, we showed the relation between the Debye potential scheme and a symmetry operator constructed from the CKY tensor. In this section we shall show the analogues of these constructions for massless Dirac fields.

Let u be an even shear-free spinor corresponding to a repeated principle null direction, and let f be a scalar field with opposite $\mathbb{C}^*$ -charge. Then the odd spinor $\psi'(f, u)$ given by

$$\psi'(f, u) = \hat{d}f u + \frac{1}{2} f \hat{D}u \quad (48)$$

is a $\mathbb{C}^*$ -scalar. The action of the Dirac operator on $\psi'(f, u)$ is

$$D\psi'(f, u) = e^a \left(\hat{\nabla}_{X_a} \hat{d}f u + \hat{d}f \hat{\nabla}_{X_a} u + \frac{1}{2} \hat{\nabla}_{X_a} f \hat{D}u + \frac{1}{2} f \hat{\nabla}_{X_a} \hat{D}u \right)$$

$$\begin{aligned}
&= \hat{d}^2 f u - \hat{d}^* \hat{d} f u + e^a \hat{d} f \widehat{\nabla}_{X_a} u + \frac{1}{2} \hat{d} f \hat{D} u + \frac{1}{2} f \hat{D}^2 u \\
&= \hat{\nabla}^2 f u + \hat{d}^2 f u + \frac{1}{2} f \hat{D}^2 u + 2 \left(\widehat{\nabla}_{\text{grad} f} u - \frac{1}{4} \hat{d} f \hat{D} u \right) \\
&= \left(\hat{\nabla}^2 f - \frac{1}{6} \mathcal{R} f \right) u + 2\mu f u + \hat{d}^2 f u
\end{aligned}$$

by the shear-free equation (13) and its integrability condition (31a). If the $\mathbb{C}^*$ -charge of u is q , then the charge of f is $-q$ and we have

$$\hat{d}^2 f = -q f \mathcal{F}$$

and so

$$\hat{d}^2 f u = -3\mu f u \quad \text{by (5.34).}$$

Thus

$$D\psi'(f, u) = \left(\hat{\nabla}^2 f - \frac{1}{6} \mathcal{R} f \right) u - \mu f u,$$

and so $\psi'(f, u)$ satisfies the massless Dirac equation if f satisfies the ‘Debye potential’ equation

$$\hat{\nabla}^2 f - \frac{1}{6} \mathcal{R} f = \frac{1}{4} \lambda_\phi f \tag{49}$$

where $\phi = u \otimes \bar{u}$ is an eigenvector of C with eigenvalue λ_ϕ . This equation is of the same form as that for the Debye potential used to construct a Maxwell field from ϕ . However, here the $\mathbb{C}^*$ -charge of the scalar potential is half that required in the Maxwell case, and the eigenvalue is also different. In the special case in which the ‘gauge terms’ are zero the shear-free spinor satisfies the twistor equation. In this case the construction of a massless Dirac solution from a scalar field satisfying the conformally-covariant wave equation is an example of what Penrose has called ‘spin raising’ [13].

As in the Maxwell case, we may form a scalar potential from a shear-free spinor and a massless Dirac solution. In the special case in which the shear-free equation reduces to the twistor equation this corresponds to Penrose’s ‘spin lowering’ [13]. Out of the shear-free spinor u and massless Dirac solution ψ we form a scalar $f(u, \psi)$ from the scalar product (u, ψ) ,

$$f(u, \psi) = (u, \psi). \tag{50}$$

(Unfortunately the notation doesn’t work well for us here. The brackets on the left-hand side denote that the scalar is constructed from u and ψ , whereas the bracket on the right-hand side denotes the symplectic product of the two spinors.) The scalar $f(u, \psi)$ has the same $\mathbb{C}^*$ -charge as u , and we may evaluate the gauged Laplacian of it by noting that the gauged covariant derivative is compatible with the spinor product:

$$\hat{\nabla}^2 f(u, \psi) = (\hat{\nabla}^2 u, \psi) + 2(\widehat{\nabla}_{X^a} u, \nabla_{X_a} \psi) + (u, \nabla^2 \psi).$$

Since u satisfies the shear-free equation (13)

$$\begin{aligned}(\hat{\nabla}_{X^a} u, \nabla_{X_a} \psi) &= \frac{1}{4}(e^a \hat{D}u, \nabla_{X_a} \psi) \\ &= \frac{1}{4}(\hat{D}u, D\psi) = 0\end{aligned}$$

since ψ satisfies the massless Dirac equation. We may then use (23) to relate the spinor Laplacian to the square of the Dirac operator, which gives zero when acting on ψ , to give

$$\begin{aligned}(\hat{\nabla}_{X^a} u, \nabla_{X_a} \psi) &= (\hat{\nabla}^2 u + \frac{1}{4}\mathcal{R}u, \psi) \\ &= (\mu u + \frac{1}{6}\mathcal{R}u, \psi)\end{aligned}$$

by the integrability condition (31b),

$$= (\mu + \frac{1}{6}\mathcal{R})f(u, \psi).$$

Thus the scalar $f(u, \psi)$ satisfies the ‘Debye potential’ equation (49). Note, however, that $f(u, \psi)$ has the same ‘charge’ as u , opposite to that required to combine with u to make a massless Dirac solution.

To construct one Dirac solution from another we need a pair of shear-free spinors with opposite ‘charges’, and this is just the case in which we have a CKY tensor. In that case we may proceed as in the Maxwell case and define the symmetry operator $\mathcal{L}_{u_1 u_2}$ by

$$\begin{aligned}\mathcal{L}_{u_1 u_2} \psi &= \psi'((u_1, \psi), u_2) \\ &= \hat{d}(u_1, \psi)u_2 + \frac{1}{2}(u_1, \psi)\hat{D}u_2.\end{aligned}\tag{51}$$

By interchanging the two spinors in this construction we could have formed the operator $\mathcal{L}_{u_2 u_1}$. However, as we shall see, these two operators are in fact the same when acting on massless Dirac fields. Since $\hat{\nabla}$ is compatible with the spinor inner product,

$$\begin{aligned}\hat{d}(u_1, \psi) &= (\hat{\nabla}_{X_a} u_1, \psi)e^a + (u_1, \nabla_{X_a} \psi)e^a \\ &= \frac{1}{4}(e_a \hat{D}u_1, \psi)e^a + (u_1, \nabla_{X_a} \psi)e^a \quad \text{by (13)}.\end{aligned}$$

Thus

$$\begin{aligned}\hat{d}(u_1, \psi)u_2 &= \frac{1}{4}(e^a u_2 \otimes \overline{e_a \hat{D}u_1})(\psi) + (e^a u_2 \otimes \bar{u}_1)(\nabla_{X_a} \psi) \\ &= \frac{1}{4}e^a(u_2 \otimes \overline{\hat{D}u_1})e_a \psi + \frac{1}{2}e^a(u_2 \otimes \bar{u}_1 + u_1 \otimes \bar{u}_2)(\nabla_{X_a} \psi) \\ &\quad + \frac{1}{2}e^a(u_2 \otimes \bar{u}_1 - u_1 \otimes \bar{u}_2)(\nabla_{X_a} \psi).\end{aligned}\tag{52}$$

If ϕ is any p -form then $e^a \phi e_a = (4 - 2p)(-1)^p \phi$. So if ϕ is odd, a sum of a 1-form and a 3-form, this becomes $e^a \phi e_a = -2\phi^\xi$. We can then use (6) to rewrite the first term in (52) using

$$\frac{1}{4}e^a(u_2 \otimes \overline{\hat{D}u_1})e_a = \frac{1}{2}\hat{D}u_1 \otimes \bar{u}_2. \quad (53)$$

The last term in (52) is made up of an anti-symmetric combination of two spinors of the same parity. Thus this even form is even under ξ , and is thus a sum of a 0-form and a 4-form. These forms commute and anticommute respectively with the 1-form e^a to enable this last term to be written in terms of the Dirac operator on ψ , which vanishes since ψ is assumed to satisfy the massless Dirac equation. Thus we can write the symmetry operator on ψ as

$$\mathcal{L}_{u_1 u_2} \psi = \frac{1}{2}(\hat{D}u_1 \otimes \bar{u}_2 + \hat{D}u_2 \otimes \bar{u}_1)(\psi) + \frac{1}{2}e^a(u_2 \otimes \bar{u}_1 + u_1 \otimes \bar{u}_2)(\nabla_{X_a} \psi).$$

In this form the two spinors enter symmetrically. Thus

$$\mathcal{L}_{u_1 u_2} \psi = \mathcal{L}_{u_2 u_1} \psi$$

for ψ a massless Dirac solution. The symmetric tensor product of the two spinors is the CKY tensor ω , in terms of which the symmetry operator can be written more concisely. Differentiating ω , and using the shear-free spinor equation, produces

$$\nabla_{X_a} \omega = \frac{1}{2}\hat{\nabla}_{X_a} u_1 \otimes \bar{u}_2 + \frac{1}{4}u_1 \otimes e_a \overline{\hat{D}u_2} + \frac{1}{2}\hat{\nabla}_{X_a} u_2 \otimes \bar{u}_1 + \frac{1}{4}u_2 \otimes e_a \overline{\hat{D}u_1}.$$

Clifford multiplication by e^a then shows that

$$d\omega - d^* \omega = \frac{3}{4}(\hat{D}u_1 \otimes \bar{u}_2 + \hat{D}u_2 \otimes \bar{u}_1),$$

by (53), and hence we may express the symmetry operator on ψ as

$$\mathcal{L}_{u_1 u_2} \psi = e^a \omega \nabla_{X_a} \psi + \frac{2}{3}d\omega \psi - \frac{2}{3}d^* \omega \psi.$$

This operator, constructed out of a self-dual ω , maps even solutions of the massless Dirac equation into odd solutions, and annihilates odd spinors. Clearly we could have taken an anti-self-dual CKY tensor and constructed a symmetry operator that maps odd Dirac to solutions to even ones. So more generally we have a symmetry operator $\mathcal{L}_\omega$ constructed out of any CKY tensor ω (with no assumptions of self-duality)

$$\mathcal{L}_\omega \psi = e^a \omega \nabla_{X_a} \psi + \frac{2}{3}d\omega \psi - \frac{2}{3}d^* \omega \psi.$$

It can be seen directly from the CKY equation, and its integrability conditions, that this is indeed a symmetry operator. In fact one can show that

$$[D, \mathcal{L}_\omega] = \left(\omega D - \frac{1}{3}d\omega + d^* \omega \right) D.$$

Operators, such as $\mathcal{L}_\omega$, whose commutator with D is of the form RD (where R is another operator) are called *R-commuting*.

Kamran and McLenaghan [19] have obtained the most general first-order *R*-commuting operator for the Dirac operator. They showed that the non-trivial terms in this operator are constructed from conformal Killing-Yano tensors of degree 1, 2 and 3. The operator constructed from the conformal Killing vector is just the Lie derivative with the appropriate conformal weight, corresponding to the well known conformal covariance of the equation. A conformal Killing-Yano 3-form is just the dual of a conformal Killing 1-form. Although they don't explicitly point it out, the operator that Kamran and McLenaghan construct from the 3-form is essentially just the operator formed from the corresponding conformal Killing vector (the operators differ by a term involving the Dirac operator and an overall factor of z). That part of their operator, $\mathcal{K}_\omega$, constructed from 'a conformal generalisation of a Penrose-Floyd tensor', is obtained from a slight modification to $\mathcal{L}_\omega$:

$$\mathcal{K}_\omega = z(\mathcal{L}_\omega - \omega D)$$

where z is the volume form. The operator $\mathcal{K}_\omega$, expressed in terms of a Killing spinor, was obtained independently by Torres del Castillo [11]. Clearly $\mathcal{L}_\omega$ and $\mathcal{K}_\omega$ only differ by a factor of i when acting on massless Dirac solutions, but the commutator of $\mathcal{K}_\omega$ with the Dirac operator becomes

$$[D, \mathcal{K}_\omega] = -\frac{2}{3}z d^* \omega D.$$

The conformal Killing-Yano tensor ω is a Killing-Yano tensor when $d^* \omega = 0$. Thus in this case the above commutator shows that $\mathcal{K}_\omega$ is a symmetry operator for the *massive* Dirac equation. In this case $\mathcal{K}_\omega$ has been interpreted as a generalised total angular momentum operator by Carter and McLenaghan [28].

VIII. Discussion and Conclusions.

There has been much work done on Debye potential methods for solving massless field equations, and on the construction of symmetry operators for these equations in algebraically-special space-times. However, the relation between different approaches has not always been made clear. We believe that the basis-independent formalism that we have given here makes it clearer to see the ingredients that have gone into the various constructions. We have explicitly shown the relationship between symmetry operators constructed from Debye potentials and those given by Kalnins, McLenaghan and Williams [12] constructed from a 4-index Killing spinor.

In this paper we have only considered the case of a four-dimensional Lorentzian space-time. One potential advantage of the approach that we have adopted is that it should be easier

to see which features of the results translate to different dimensions and signatures. The adapted basis of the Newman-Penrose formalism is of course optimised for the Lorentzian case, and any explicit component expression will become unwieldy in higher dimensions. Whereas one can always (subject to the usual topological caveats) introduce spinors in any number of dimensions, there are many features of the 2-spinor calculus that are special to four dimensions. Four dimensions are also rather special for Maxwell's equations: 2-forms are 'middle forms'. If we regard Maxwell's theory as a gauge theory then the Maxwell forms will be 2-forms regardless of the dimension. However, the geometrical relationship between null Maxwell solutions and shear-free null geodesics is carried over to higher dimensions to a relationship between 'middle forms' and certain foliations [29]. Conformal Killing-Yano tensors are of course readily introduced in any number of dimensions (indeed Tachibana [2, 30] considered arbitrary dimensions, although positive-definite signature). In four dimensions conformal Killing-Yano 2-forms are the only non-trivial generalisation of conformal Killing vectors. The 0-forms and 4-forms are parallel if they are CKY whilst the 3-form is just the Hodge dual of a conformal Killing vector. In higher dimensions there are of course more possibilities. We hope to see which of these tensors can be used to generate symmetry operators for various massless field equations.

The symmetry operator given by the Debye scheme is expressed in terms of a '4-index Killing spinor' that is formed from the tensor product of the CKY 2-form. However, for this to be a symmetry operator we only need the '4-index Killing spinor' equation to be satisfied; it is not necessary that the '4-index Killing spinor' be a product of '2-index' ones. At this point in the paper we have not given full details of how our calculations were performed. We hope to give a fuller account of 'generalised conformal Killing-Yano tensors' in a more general setting later.

There are a number of aspects of Debye potentials and symmetry operators that we have not discussed here. One is the application to higher spin fields. In conformally-flat space-times Debye potentials can readily be extended to include massless fields of arbitrary spin [13]. A Debye potential prescription for spin- $\frac{3}{2}$ fields in electro-vac space-times, within the generalised Goldberg-Sachs class, has been given by Torres del Castillo, who has also discussed Debye potentials for spin-2 fields [31, 32, 8]. An important application of symmetry operators, and their relation to CKY tensors, is the question of separation of variables. The separation constants obtained in this procedure can be given an intrinsic characterisation in terms of eigenvalues of symmetry operators. Torres del Castillo [33] has shown how, for Maxwell fields, the Starobinsky constant is given by the symmetry operator obtained via Debye potential methods in the Plebański-Demiański background and Silva-Ortigoza [34] has presented a similar analysis for the Rarita-Schwinger (spin- $\frac{3}{2}$) equation. Recently Kalnins, Williams and Miller [35] have given a detailed account of the separation of variables for electromagnetic and gravitational perturbations in the Kerr space-time using Hertz potentials.

As a final note, we point out that not only have Debye potential methods been successfully applied to many cosmologically interesting space-times [36, 37], they have also lead to new solutions in the seemingly unrelated fields of isotropic elastic media [38] and force-free magnetic fields [39].

Acknowledgment. IMB thanks R. W. Tucker for kindling his interest in this area and for hospitality whilst visiting the University of Lancaster. This research was supported by a Newcastle University research management committee grant. PRC was supported by an APRA scholarship.

References

- [1] J M Cohen and L S Kegeles. Electromagnetic fields in curved spaces: A constructive procedure. *Physical Review D*, 10(4):1070–1084, 1974.
- [2] Shun ichi Tachibana. On conformal Killing tensor in a Riemannian space. *Tôhoku Math. Journ.*, 21:56–64, 1969.
- [3] I Robinson. Null electromagnetic fields. *Journal of Mathematical Physics*, 2(3):290–291, 1961.
- [4] W Dietz and R Rüdiger. Shearfree congruences of null geodesics and Killing tensors. *Gravitation and General Relativity*, 12(7):545–562, 1980.
- [5] L S Kegeles and J M Cohen. Constructive procedure for perturbations of space-times. *Physical Review D*, 19(6):1641–1664, 1979.
- [6] J M Stewart. Hertz-Bromwich-Debye-Whittaker-Penrose potentials in general relativity. *Proceedings of the Royal Society of London A*, 367:527–538, 1979.
- [7] G F Torres del Castillo. Null Strings and Hertz potentials. *Journal of Mathematical Physics*, 25(2):342–346, 1984.
- [8] R M Wald. Construction of solutions of gravitational, electromagnetic, or other perturbation equations from solutions of decoupled equations. *Physical Review Letters*, 41(4):203–206, 1978.
- [9] S A Teukolsky. Perturbations of a rotating black hole. I. Fundamental equations for the gravitational, electromagnetic, and neutrino-field perturbations. *The Astrophysical Journal*, 185:635–647, 1973.
- [10] G F Torres del Castillo. The Teukolski-Starobinsky identities in type D vacuum backgrounds with cosmological constant. *Journal of Mathematical Physics*, 29(9):2078–2083, 1988.
- [11] G F Torres del Castillo. Killing spinors and massless spinor fields. *Proceedings of the Royal Society of London A*, 400:119–126, 1985.

- [12] E G Kalnins, R G McLenaghan and G C Williams. Symmetry operators for Maxwell's equations on curved space-time. *Proceedings of the Royal Society of London A*, 439:103–113, 1992.
- [13] R Penrose and W Rindler. *Spinors and Space-time*, volume 2. Cambridge University Press, Cambridge, 1986.
- [14] I M Benn and R W Tucker. *An Introduction to Spinors and Geometry with Applications in Physics*. IOP Publishing Ltd, Bristol, 1987.
- [15] J A Thorpe. Curvature and Petrov canonical forms. *Journal of Mathematical Physics*, 10(1):1–7, 1969.
- [16] B O'Neill. *The Geometry of Kerr Black Holes*. A K Peters, Wellesley, Massachusetts, 1995.
- [17] Kentaro Yano. Some remarks on tensor fields and curvature. *Annals of Mathematics*, 55(2):328–347, 1952.
- [18] M Walker and R Penrose. On quadratic first integrals of the geodesic equations for the type {22} space-times. *Communications in Mathematical Physics*, 18:265–274, 1970.
- [19] N Kamran and R G McLenaghan. Symmetry operators for the neutrino and Dirac fields on curved space-time. *Physical Review D*, 30(2):357–362, 1984.
- [20] I M Benn. A unified description of null and non-null shear-free congruences. *Journal of Mathematical Physics*, 35(4):1796–1802, 1994.
- [21] P Sommers. Properties of shear-free congruences of null geodesics. *Proceedings of the Royal Society of London A*, 349:309–318, 1976.
- [22] J N Goldberg and R K Sachs. A theorem on Petrov types. *Acta Physica Polonica*, 12:13–23, 1962.
- [23] E Newman and R Penrose. An approach to gravitational radiation by the method of spin coefficients. *Journal of Mathematical Physics*, 3(3):566–578, 1962.

- [24] I Robinson and A Schild. Generalization of a theorem by Goldberg and Sachs. *Journal of Mathematical Physics*, 4(4):484–489, 1963.
- [25] O Laporte and G E Uhlenbeck, *Physical Review*, 37:1380, 1931.
- [26] A Nisbit. Hertzian electromagnetic potentials and associated gauge transformations. *Proceedings of the Royal Society of London A*, 250:250–263, 1955.
- [27] E Mustafa and J M Cohen. Hertz and Debye potentials and electromagnetic fields in general relativity. *Classical and Quantum Gravity*, 4:1623–1631, 1987.
- [28] B Carter and R G McLenaghan. Generalized total angular momentum operator for the Dirac equation in curved space-time. *Physical Review D*, 19(4):1093–1097, 1979.
- [29] L P Hughston and L J Mason. A generalised Kerr-Robinson theorem. *Classical and Quantum Gravity*, 5:275–285, 1988.
- [30] Shun ichi Tachibana and Toyoko Kashiwada. On the integrability of Killing-Yano’s equation. *Journal of the Mathematical Society of Japan*, 21(2):259–265, 1969.
- [31] G F Torres del Castillo. Gravitational perturbations of algebraically special space-times via the $\mathcal{HH}$ equation. *Journal of Mathematical Physics*, 27(6):1586–1591, 1986.
- [32] G F Torres del Castillo. Perturbations of solutions of the Einstein-Maxwell equations with a null background electromagnetic field. *Journal of Mathematical Physics*, 37(8):4053–4061, 1996.
- [33] G F Torres del Castillo. The separability of Maxwell’s equations in type-D backgrounds. *Journal of Mathematical Physics*, 29(4):971–977, 1988.
- [34] G Silva-Ortigoza. Killing spinors and separability of Rarita-Schwinger’s equation in type $\{2, 2\}$ backgrounds. *Journal of Mathematical Physics*, 36(12):6929–6936, 1995.
- [35] E G Kalnins, G C Williams and W Miller Jr. Intrinsic characterization of the separation constant for spin one and gravitational perturbations in Kerr geometry. *Proceedings of the Royal Society of London A*, 452:997–1006, (1996).

- [36] S V Dhurandhar, C V Vishveshwara and J M Cohen. Electromagnetic fields in space-times with local rotational symmetry. *Physical Review D*, 21(10):2794–2804, 1980.
- [37] S V Dhurandhar, C V Vishveshwara and J M Cohen. Neutrinos in perfect fluid space-times with local rotational symmetry. *Physical Review D*, 26(10):2598–2610, 1982.
- [38] G F Torres del Castillo. Solution of the inhomogeneous equations of equilibrium for an isotropic elastic medium. *Revista Mexicana de Física*, 41(5):695–702, 1995.
- [39] I M Benn and Jonathan Kress. Force-free fields from Hertz potentials. *Journal of Physics A:Mathematical and General*, 29:6295–6304, (1996)